

# Studia Orchidacearum Japonicarum VI\*

## Orchidaceae Formosae Et Insulae Kōtōsyo Novae Atque Criticae

auctore

Noriaki Fukuyama

Received April 24, 1935

### 48) **Orchis Kunihikoana** MASAMUNE et FUKUYAMA, sp. nov.

*Orchis kiraishiensis* HAYATA form. *leucantha* MASAMUNE, in Journ. Trop. Agr. Taihoku Imp. Univ. III. (1931) p. 246.

*Orchis kiraishiensis* HAYATA var. *leucantha* MASAMUNE, in Trop. Hort. III. (1933) p. 45.

Terrestris, erecta, glaberrima, c. 10 cm. alta, tuberibus ovoideis 5–10 mm. longis pilosiusculis, radicibus paucis brevibus filiformibus flexuosis puberulis. Caulis tenuis, teres, strictus vel plus minusve flexuosus, glaber, ad medium subconferte vel remote bifoliatus, interdum unifoliatus, dimidio inferiore paucivaginatus, vaginis membranaceis glabris caulem complete amplectentibus apice truncatis vel obtusis acutisve.

Folia suberecta, patentia, oblanceolata vel linearis-oblonga vel lanceolata, apice acuta vel breve acuta, basi longe decurrentia et vaginaria, lamina 3–4 cm. longa, medio 5–10 mm. lata, margine integra, in exsiccatum membranacea, utrinque glabra.

Flores 2–5, in apice caulis subdense racemosi, subsecundi, albescentes, bracteis foliaceis lanceolatis apice acuminatis basi paullo contractis utrinque glabris florem paullo vel diplo superantibus, pedicellis cum ovariis c. 5–10 mm. longis rectis glabris apice basique contractis. Sepalum intermedium concavulum, oblongum, apice acutiusculum, basi vix contractum, margine integrum, c. 6.5 mm. longum, medio 2.5 mm. latum, glabrum, membranaceum, trinervatum; sepala lateralia erecto-patentia, ovato-oblonga, obliqua, latere superiore truncata, apice acutiuscula vel apiculata, basi inaequilaterale contracta, c. 8 mm. longa, infra medium 3.5 mm. lata, glabra, membranacea, trinervata. Petala cum sepallo intermedio in galeam coninventia, ovato-oblonga, valde obliqua, apice obtusiuscula, basi vix inaequilaterale contracta, sepallo intermedio subaequilonga, membranacea, glabra, trinervata.

Labellum erecto-patens, e basi perlate cuneatum, supra perbrevissime

\* Continued from Bot. Mag. (Tokyo), 49 (1935), p. 444.

puberulum, subtus glabrum, supra medium trilobatum, inter apices loborum lateralium explanatorum c. 1.2 cm. latum, lobis lateralibus obliquis apice obtusis vel truncato-rotundatis margine irregularibus, lobo intermedio quam lateralibus longiore quadrangulato antice retuso vel subbilobulato-emarginato ad centrum breve aristato infra apicem c. 6 mm. lato margine irregulari, calcare cylindrico recto apice obtuso ad basin plus minusve ampliato pedicellis paullo breviore medio c. 1 mm. diametente. Columna c. 3 mm. alta.

Nom. Jap. Kumoma-tidori (nov.)

Hab. Formosa; praef. Taityū, in herbidis alpinis monte Taisuikutsan, (D. ORIMOTO) Jul. 1934. no. 4704. Typus in Herb. Orch. Fuk.)

49) **Amitostigma alpestre** FUKUYAMA, sp. nov.

Terrestre, pusillum, erectum, glaberrimum, 5–10 cm. altum, tuberibus ovoideis vel globosis 5–10 mm. longis pilosiusculis. Caulis tenuis, teres, substrictus vel plus minusve flexuosus, glaber, dimidio inferiore laxe 1-2-foliatus, ima basi bivaginatus, vaginis membranaceis blabris amplexieaulibus apice truncatis vel acutis.

Folia suberecta, patentia, oblanceolata vel anguste oblonga, apice breve acuta, basi longe decurrentia et amplectentia, c. 5 cm. longa, supra medium 5–10 mm. lata, margine integra, in exsiccato membranacea, utrinque glabra.

Flores 2–3 raro 4, in apice caulis sublaxe dispositi, subsecundi, lilacini vel lilacino-punctati, subapertientes, bracteis foliaceis lanceolatis apice acuminatis basi contractis utrinque glabris inferioribus florem superantibus superioribus sensim brevioribus, pedicellis cum ovariis c. 5–10 mm. longis rectis glabris apice basique contractis. Sepalum intermedium concavulum. ovato-oblongum, apice obtusiuseulum, basi paullo contractum, margine integrum, 4–5 mm. longum, infra medium vix 2 mm. latum, glabrum. membranaceum, trinervatum; sepala lateralia ovato-oblonga, paullo obliqua. quam sepalum intermedium paullo majora, apice acutiuscula vel obtusiuscula. basi contracta, glabra, membranacea, trinervata. Petala cum sepalo intermedio in galeam conniventia, ovato-lanceolata, obliqua, apice acutiuscula, basi contracta, margine integra, c. 4 mm. longa, supra basin c. 1 $\frac{1}{2}$  mm. lata. utrinque glabra vel perbrevissime puberula, membranacea, 1-3-nervata.

Labellum erecto-patens, supra perbrevissime puberulum, subtus glabrum, membranaceum, c. 6 mm. longum. medio trilobatum, inter apices loborum lateralium explanatorum c. 5 mm. latum, ad basin cuneato-attenuatum, lobis lateralibus oblique oblongis apice acutis vel obtusis vel truncatis margine integris vel leviter erosis latere exteriore c. 4 mm. interiore c. 2 mm. longis a lobo intermedio angulo 45° egressis, lobo intermedio quam lateralibus longiore obovato-quadrangulato apice truncato ad summum leviter