

# Studia Orchidacearum Japonicarum VII\*

## Orchidaceae Formosanae Novae Atque Criticae

auetore

Noriaki Fukuyama

Received April 2, 1935

### 57) *Cymbidium Syunitianum* FUKUYAMA, sp. nov.

Terrestre, erectum, 40–66 cm. altum, caulescens, rhizomate valde abbreviato, radicibus incrassatis teretibus c. 4–7 mm. diametentibus elongatis rectiusculis rigidis in exsiccato rugulosis, caulis elongato-cylindraceis omnibus vaginis acuminatis nervosis amplectentibus.

Folia c. 4, ad apicem caulis alternatim fasciculata, erecto-patentia, oblanceolata, apice acuta vel acuminata, basi ad petiolum longe decurrentia, margine integra, tenuiter coriacea, lamina 20–30 cm. longa, supra medium 4–5 cm. lata, utrinque glabra, 3-5-nervata, ad nervos subtus elevata supra laevia, inter nervos multi-venulosa.

Scapus e basi caulis oriundus, simplex, strictus, glaber, viridescens, prope basin pedicellorum violaceus, c. 25 cm. longus, superiore sublaxe c. 10-floratus, inferiore pauci-vaginatus, bracteis lanceolato-linearibus apice acuminatissimis basi non contractis inferioribus c. 2.5 cm. longis superioribus sensim minoribus viridescentibus glabris multi-nervatis.

Flores apertentes, viridescentes, glabri, haud odorati, pedicellis eum ovarii rectis vel curvatis glabris viridescentibus quam bracteis paullo vel diplo longioribus. Sepala patentia, glabra, in vivo viridescentia, rigidiuscula, aequilonga, aequilata, subaequiformia, 5-nervata, sepalo intermedio linearis-oblongo apice acuto ad basin vix attenuato c. 3 cm. longo medio c. 6 mm. lato, sepalis lateralibus intermedio similibus sed plus minusve obliquis. Petala columnam solete amplectentia, recta, oblique ligulata, ad summam acutiuscula, basi paulo contracta, margine integra, quam sepala paulo breviora latioraque, viridescentia, intus medio violaceo-lineata, glabrata.

Labellum quam petala paulo brevius, columnam amplectens, eircuitu ovato-lanceolatum, infra medium paulo curvatum, anteriore rectum, basi rotundatum, e medio apicem versus sensim caudato-angustatum, parte caudata margine inaequilaterale revolutum, ad summam acutum, infra

\* Continued from Bot. Mag. (Tokyo), 49 (1935), 444.

20 cm. longis vivo 1–1.5 mm. diametentibus sublaxe et distiche foliatis cum vaginis foliorum toto facie obtectis sparse et minute purpureo-puncticulatis glabratris.

Folia patentia, alterna, disticha, deorsum paullo arcuata, anguste oblonga, apice oblique breveque acuta, basin versus attenuata, in petiolos breves abeuntia, 1.5–3 em. longa, medio c. 5 mm. lata, crassa, supra viridia et medio leviter unisulcata, subtus convexa pallidioraque, utrinque glabra et non maculata.

Racemus lateralis, plerumque divaricato-bifloratus, breviter pedunculatus, pedunculis rectis c. 1 em. longis basi paucis-squamulatis purpureo-maculatis glabris, bracteis minutis squamulis subaequiformibus subaequimagnisque triangulatis apice acutis basi non contractis glabratris.

Flores subapertientes, flavescentes vel viridi-flavescentes, rubro-fusco-maculati, inter apices sepalorum intermediorum et fundos saccorum c. 1 em. diametentes, pedicellis rectis bracteas multo superantibus 7–10 mm. longis viridescentibus glabris. Sepala crassa, leviter, concava, glabra, flavescentia, intus rubro-fusco-maculata, sepalo intermedio elliptico apice obtuso ad basin contracto c. 5 mm. longo medio vix 3 mm. lato, sepalis lateralibus intermedio similibus sed paullo minoribus obliquisque. Petalum crassa, leviter concava, glabra, flavescentia, intus rubro-fusco-maculata, obovato-oblonga, apice obtusa, basin versus angustata, sepalo intermedio subaequilonga et aequilata.

Labellum ad basin columnae sessile, hypochilio saccatum, sacco oblongo-hemisphaericō paullo complanato apice rotundato c. 6 mm. alto flavescente intus glabro purpureo-punctato ore purpurascenti-colorato, epichilio late triangulato apice obtuso basi truncato margine integro vel vix irregulari 3–4 mm. longo basi c. 7 mm. lato albescenti supra medio callo viridescenti-flavo purpureo-punctato instrueto ad callum puberulo subtus glabro. Columna brevis, crassa.

Nom. Jap. Kintyaku-ran (nov.)

Hab. Formosa: praef. Taihoku, in *Sakakietis* dorso montis Haro, c. 1000 m. alt., (SUZUKI-T. Mart. 20, 1935. no. 4715. Typus in Herb. Oreh. Fuk.)

### 65) **Gastrochilus rupestris** FUKUYAMA, sp. nov.

Epiphyticus, adpresso prostratus, radicibus e caule emittentibus filiformibus vel vermiformibus leviter flexuosis 0.5–1.5 mm. diametentibus glabris, caulinis 5–10 em. longis vivo 1–1.5 mm. diametentibus subdense foliatis cum vaginis foliorum toto facie obtectis minute purpureo-puncticulatis glabratris.

Folia patentia, alterna, disticha, parva, elliptica, apice breve acuta.

ad basin contracta, subito in vaginas abeuntia, 8–10 mm. longa, medio 3.5–6 mm. lata, crassa, rigida, supra leviter concava et medio unisulcata, subtus convexa, utrinque purpureo-punctata vel purpureo-maculata, glabrata.

Racemus lateralis, semper divaricato-bifloratus, breviter pedunculatus, pedunculis rectis clavatis c. 5 mm. longis remote pauci-squamulatis glabris, bracteis minutis squamulis subaequiformibus triangulatis apice acutiusculis basi non contractis glabratis.

Flores apertentes, viridi-flavescentes, purpureo-maculati, inter apices sepolorum intermediorum et fundos saceorum c. 1 cm. diametentes, pedicellis paulo curvatis bracteas multo superantibus 5–8 mm. longis glabratiss. Sepala crassa, incurva, concava, utrinque glabra, rubro-fusco-maculata, sepalo intermedio orbiculato-elliptico apice rotundato-obtuso basi contracto vix 5 mm. longo medio vix 3 mm. lato, sapalis lateralibus obovato-ellipticis paulo obliquis apice obtusis basin versus attenuatis 5–6 mm. longis medio c. 3 mm. latis. Petala crassa, leviter concava, glabra, rubro-fusco-maculata, sepalo intermedio subaequiformia, apice rotundata, basi contracta, c. 5 mm. longa, medio 3 mm. lata.

Labellum ad basin columnae sessile, hypochilio saccatum, sacco complanato-hemisphaericō apice rotundato c. 5 mm. alto intus et extus glabro, epichilio late semiorbiculato anteriore truncale-rotundato ad centrum emarginato vel cordato margine anteriore irregulariter et leviter crenulato c. 3.5 mm. longo basi c. 7.5 mm. lato supra ad medium callo instructo ad callum puberulo subtus glabro. Columna brevis, crassa.

Nom. Jap. Hisasi-kintyaku-ran (nov.)

Hab. Formosa; praef. Sintiku, Epiphytus in saxis monte Ritō-san, prope Takadai, c. 1500 m. alt., (N. FUKUYAMA) Jan. 7, 1932. no. 4716. Typus in Herb. Orch. Fuk.)

(To be continued)

Mar. 30, 1935

The Laboratory of Systematic Botany and Plantecology  
Taihoku Imperial University, Formosa, Japan