

Contribution to the Orchid Flora of the Ryûkyû  
Archipelago\* II.

By Noriaki FUKUYAMA (edited by G. MASAMUNE)

福山伯明：琉球列島のラン II (正宗嚴敬編)

6) *Thrixspermum neglectum* FUKUYAMA, sp. nov. (Sect. *Dendrocolla* Bl.)

Epiphyticum, subrectum, pumilus, c. 3-5 cm. altum, caulis abbreviatis c. 1 cm. altis teretibus cum vaginis foliorum toto facie obtectis ad basin radicantibus, radicibus elongatis flexuosis filiformibus c. 1 mm. diametentibus. Folia 3-5-7, plana, laevigata, coriacea, approximativ et alternativ disposita, angulo 45° a caule divaricata, oblonga, ad summum oblique simpliciter apiculata vel minute biapiculata, margine integra, ad basin contracta, cum vaginis equitantibus c. 4 mm. longis multinervatis articulata, lamina c. 2-5 cm. longa, medio 8 mm. lata, utrinque viridia, glabra, ad costam supra sulcata, subtus subcarinatim paululo elevata. Scapus axillaris, foliis longior vel brevior, pergracilis, substrictus, viridis, glaber, distante (1-2)-squamatus, squamis minutis deltoideis, rhachibus brevibus paulo incrassatis viridibus glabris dense bracteatis, bracteis deltoideis pedicellis multo brevioribus vix 1 mm. longis. Flores subelausi, albicans, pedicellis cum ovario subrectis gracilibus glabris c. 2.5-4 mm. longis. Sepalum intermedium album, ellipticum vel obovatum, apice rotundato-obtusum, basi angustatum, concavulum, c. 4 mm. longum, medio 2.5 mm. latum, membranaceum, glabrum; sepala lateralia alba, late elliptica, paulo obliqua, apice obtusiuscula, basi contracta, sepalum intermedium aequilonga sed latiora. Petala labe, late elliptica, paulo obliqua, apice obtusa, basi contracta, sepalo intermedio subaequilonga sed latiora. Labellum albicans, ad basin et margine anteriore intus flavidum, interdum extus prope basin flavidum-notatum, basi pede columnae articulatum, 3-lobum, c. 3 mm. altum, inter apices loborum lateralium c. 7 mm. latum, intus prope apicem callis duobus globosis dense pilosis luteis instructum, partibus laterilibus exterioribus callorum pr. cessibus linearibus dispositum, infra callos lamellis luteis vel flavidis instructum, lobis lateralibus erectis subfalcato-ligulatis apice crenulatis, lobo intermedio obsoleto semiorbiculato vel deltoideo interdum plus minusve denticulato. Columna brevis, subrecta, c. 1 mm. longa, basi ad pedem c. 2 mm. longum producta. Anthera orbiculato-cucullata. Nom. Jap. Hagukure-nagamirana (FUKUYAMA nov.).

Hab. Ryûkyû. Ins. Iriomote: in laurisilvis praeter flumen Umuti-gawa, ad Hutamata (N. FUKUYAMA, Jun. 5, 1938, n. 7448). Formosa. Praef. Taihoku: in laurisilvis praeter flumen Tôgo-kei, ad Agyoku (N. FUKUYAMA, Dec. 28, 1935. Colitur in horto et floret in mense Mai., 1936, n. 7447-Typus in Herb. Nation. Taiwan Univ.).

\* Contribution from the Laboratory of the Systematic Botany, Faculty of Science, Kanazawa University, No. 10.

Distrib. Formosa, Ryūkyū.

Near to *Thrixspermum kusukusensis* HAYATA, but distinguishable from it by the broader sepals and petals and the denticulate side lobes of the lip.

7) **Staurochilus luchuensis** FUKUYAMA, comb. nov.—*Stauropsis luchuensis* ROLFE, in Kew Bull. (1907) 131.—*Pandopsis luchuensis* SCHLECHTER, in FEDDE Repert. X, 196 (1911).—*Cleisostoma ionosimum* LINDEL. form. *lutchuense* MAKINO, in Tokyo Bot. Mag. XXI, 60 (1907).

Nom. Jap. Nyūmenran, Iriomoteran.

Hab. Ryūkyū. Ins. Isigaki: in silvis montis Omoto-dake. (sive E. TAKAMINE olim.).

Distrib. Ryūkyū australis.

8) **Cheirostylis liukiuensis** MASAMUNE, in Journ. Soc. Trop. Agr. Taihoku Imp. Univ. Formosa, II, 36 (1930).—*Cheirostylis Kanashiroi* OHWI, in Act. Phytotax. et Geobot. VII, 134 (1938).

Nom. Jap. Akaba-susuran (MASAMUNE), Okumaran (OHWI).

Hab. Ryūkyū. Ins. Uchina: Kunigami (S. SONOHARA, typus in Herb. Nation. Taiwan Univ.); in monte Katu'u-dake (Y. TAIRA, Mart. 30, 1939, n. 492.); in monte Kūrā (Y. TAIRA, Mart. 6, 1938, n. 222.).

The original description of *C. Kanashiroi* OHWI is satisfactory coincide with the type specimen of *C. liukiuensis* MASAMUNE preserved in our herbarium. I consider that these species may be conspecific, in which case *C. liukiuensis* MASAMUNE would by priority become the name of the species.

9) **Calanthe amamiana** FUKUYAMA, sp. nov. (Sect. *Calothrysus*.).

Terrestris, ecaulescens, c. 30 cm. alta; radices filiformes, elongatae, leviter flexuosa, haud ramosae, sicco siccatae, glabrae vel villosulae. Folia vulgo 1-2, basilaria, suberecta, elliptica, apice acuta, margine integra, basi in petiolum gracilem abeuntia; lamina c. 15-20 cm. longa, medio fere 5-6 cm. lata, sicco membranacea, utrinque glabra, subtus prominente 5-nervia, inter nervos venulosa; petiolus quam lamina plus minusve brevior vel subaequans, strictus, extus striato-nervosus, involutus, glaber; folia juniora basi vaginis amplectentibus paucis instructa. Scapus ex fasciculo foliorum oriundus, foliis subaequilongus, strictus, gracilis, teres, pubescens, distante pauci-squamatus, superiore laxe fere 4-lorus; bracteae persistentes, ovato-lanceolatae, acuminatae, c. 5 mm. longae, glabrae. Flores in genere mediocres, color mihi nondum notus; pedicellus cum ovario erecto-patens, rectus vel arcuatus, c. 15 mm. longus, puberulus. Sepalum intermedium oblongo-ellipticum, apice acutum, ad basin angustatum, c. 17 mm. longum, membranaceum, glabrum 5-nervium; sepala lateralia intermedio similia sed paullo obliqua. Petala linearia, lente falcata, apice acuta, basin versus vix attenuata, sepalis paullo breviora, membranacea, glabra, 3-nervia. Labellum glabrum, calearatum; limbus c. 10 mm. longus, c. 12 mm. latus, infra medium 3-lobus; lobus intermedius oblongus, 6 mm. longus, medio c. 2.5 mm. latus, apicem versus plus minusve dilatatus, apice submarginatus cum

apiculo interjectus, medio lamellis 3 inconspicuis parallelis ornatus; lobi laterales intermedio majores, oblique quadrato-obovati, c. 4 mm. lati; calcar paullo arcuatum, cylindricum, apice subacute, glabrum, c. 1 cm. longum. Columna brevis, crassa, antice adscendens.

Nom. Jap. Amami-ebine (FUKUYAMA nov.).

Hab. Ryūkyū. Ins. Oshima: sine loco speciali (T. KAMIVA, Apr. 7, 1910, n. 7443-Typus in Nation. Taiwan Univ.).

It resembles *Calanthe nipponica* MAKINO, but the details of the lip are characteristic. The side lobes are much broader than the inner one, the spur is twice as long.

10) **Goodyera Sonoharae** FUKUYAMA, sp. nov.

Terrestris, erecta, pusilla, 10-15 cm. alta, rhizomate decumbente caulinormi tereti viride glabro laevi ad nodos radicante, radicibus vermiformibus villosis. Caulis simplex, erectus, teres, glaber, basi laxe sunsum subconferte (5-8)-foliatus, supra folia vaginulis paucis erectis acutis obsessus, inflorescentiam versus dense glandulosopilosus. Folia erecto-patentia, ovata vel ovato-oblonga, apice apiculata, basi contracta, subito ad petiolum abeuntia, lamina 2-3 cm. longa, infra medium 1-1.7 cm. lata, margine integra, vivo undulata, supra subsericeo-laevigata, toto viridia, utrinque glabra, papyracea, 5-nervia, inter nervos reticulato-venulosa, sunsum in vaginulae decessentia, petiolis 1-1.2 cm. longis glabris basi dilatata vaginantibus. Inflorescentia subconferte multi flora, parte florifera c. 3-5 cm. longa; rachis brevis; bracteis suberectio lanceolatis subulato-acuminatis dorso hirsutis pedicellis longitudine paullo superantibus. Flores subcausi, vide collectoris pallide olivaceo-rosei, extus hirsuti, pedicellis rectis teretibus dense glandulosopilosus c. 7 mm. longis. Sepalum intermedium naviculariforme concavum, oblongum, apice obtusum, basi paullo constrictum, c. 6.5 mm. longum, infra medium explanata 3-3.5 mm. latum, extus breve hirsutum, intus glabrum, membranaceum, tri-nervium. Sepala lateralia ovato-oblonga, paullo obliqua, apice obtusa, basi vix contracta, sepalo intermedio subaequali longa vel breviora, c. 1 mm. lata, extus breye hirsuta, intus glabra, membranacea, tri-nervia. Petala oblique semirhomboidea, apice acutiuscula, basi cuneatungustata, c. 6 mm. longa, medio 2 mm. lata, utrinque glabra, membranacea, 1-nervia. Labelum crassiusculum, marginem versus membranaceum, subrectum, globosecucullatum, antice in lobulum brevem obtusiusculum productum, c. 4.5-5 mm. longum, c. 2-2½ mm. altum, extus glabrum, intus pilis longis sparsim instructum.

Nom. Jap. Kunigami-syusuran (nov.).

Hab. Ryūkyū. Ins. Okinawa-hontō: in laurisilvis inter Hentona et Aha (S. SONOHARA, Sept. 1935, n. 7017-Typus in Herb. Nat. Taiwan Univ.); in laurisilvis montis Tainu-take (N. FUKUYAMA, Sept. 1, 1938, n. 7448).

Near to *Goodyera morrisonicola* HAYATA, but distinguishable from it by the smaller hirsute flowers and differently shaped lips.

11) **Goodyera commelinoides** FUKUYAMA, sp. nov.